



REVIEW

Air pollution, oxidative stress and dietary supplementation: a review

I. Romieu*, F. Castro-Giner[#], N. Kunzli^{#,†} and J. Sunyer^{#,†}

ABSTRACT: The aim of the present review was to provide an up-to-date overview of the biological and epidemiological evidence of the role of oxidative stress as a major underlying feature of the toxic effect of air pollutants, and the potential role of dietary supplementation in enhancing antioxidant defences.

A bibliographic search was conducted through PubMed. The keywords used in the search were “air pollutant”, “oxidative stress”, “inflammation”, “antioxidant polyunsaturated fatty acids” and “genetics”. In addition, the authors also searched for biomarkers of oxidative stress and nutrients.

The review presents the most recent data on: the biological and epidemiological evidence of the oxidative stress response to air pollutants; the role of dietary supplementation as a modulator of these effects; and factors of inter-individual variation in human response. The methodology for further epidemiological studies will be discussed in order to improve the current understanding on how nutritional factors may act.

There is substantial evidence that air pollution exposure results in increased oxidative stress and that dietary supplementation may play a modulating role on the acute effect of air pollutants. Further epidemiological studies should address the impact of supplementation strategies in the prevention of air-pollution-related long-term effects in areas where people are destined to be exposed for the distant future.

KEYWORDS: Air pollution, antioxidants, nutrition, oxidative stress

Epidemiological studies have clearly shown that air pollution exposure is associated with a range of respiratory and cardiovascular health effects and increased mortality [1]. Recent research has identified oxidative stress as one potential feature underlying the toxic effect of air pollutants, which trigger a number of redox sensitive signalling pathways, such as those of inflammatory response and cytokine production [2–5]. Toxicity may arise from an imbalance of biological pro-oxidant and antioxidant processes [6] linked to increased exposure to oxidants or the presence of impaired antioxidant defences [7, 8]. This imbalance has long been recognised in investigations of ozone (O₃) [9], one of the most potent oxidants, and more recent studies have focused on this particular mechanistic hypothesis [10]. Since diet is a major source of antioxidants, it is important to examine whether antioxidant defence mechanisms could be increased by dietary means to protect against air pollutants as this could have

major public health consequences [11]. To provide an up-to-date overview on the biological and epidemiological evidence of the role of oxidative stress as a major underlying feature of the toxic effect of air pollutants and the potential role of dietary supplementation as an enhancer [11] of antioxidant defences, a bibliographic search was conducted through PubMed. The keywords used in the search were “air pollutant”, “oxidative stress”, “inflammation”, “antioxidant” (vitamin C, vitamin E, carotenoids), “polyunsaturated fatty acids” (PUFA) and “genetics”. In addition, the current authors searched for biomarkers of oxidative stress, biomarkers of antioxidant intake (selenium, flavonoids, carotenoids, vitamin C, vitamin E), and n-3 PUFA. Various recent reviews have been published on these issues [1–5, 7–10, 12–34], therefore, the present authors refer to these and mostly focus on the latest findings. Thus, the purpose of this up-to-date overview is five-fold. First, the relevance of oxidative stress as a common mechanism for

AFFILIATIONS

*Instituto Nacional de Salud Publica, Cuernavaca, Mexico.

[#]Instituto Municipal de Investigaciones Médicas, and

[†]Institut Català de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

CORRESPONDENCE

I. Romieu
Instituto Nacional de Salud Publica
Av. Universidad No. 655
Col. Santa Maria Ahuacatlan
Cuernavaca
Morelos 62508
Mexico
Fax: 52 7773111148
E-mail: iromieu@correo.insp.mx

Received:

October 02 2006

Accepted after revision:

August 20 2007

SUPPORT STATEMENT

The work was supported by the National Center for Environmental Health, Centers for Disease Control and Prevention (Atlanta, GA, USA), the GA²LEN project (EU contract FOODCT-2004-506378) and the Spanish Ministry of Education and Science (SAB2004-0192).

STATEMENT OF INTEREST

None declared.

European Respiratory Journal
Print ISSN 0903-1936
Online ISSN 1399-3003

effects of ambient air pollutants will be summarised. Secondly, the role of antioxidants in oxidative stress will be briefly discussed. Thirdly, the evidence for dietary supplements as modulating the adverse effects due to air pollution will be reviewed. Fourthly, the relevance of factors that may interact with a subjects' response to exogenous oxidative stress will be discussed. Finally, the need to further investigate the relevance of dietary supplementation as an approach to protect from adverse effects of air pollution will be discussed.

BIOLOGICAL AND EPIDEMIOLOGICAL EVIDENCE

Oxidative stress and air pollutants

Several air pollution components have been related to particulate toxicity. An important determinant of the acute inflammatory response appears to be the dose of bio-available transition metals (such as copper, vanadium, chromium, nickel, cobalt and iron), organic compounds (such as polycyclic aromatic hydrocarbons) and biological fractions (such as endotoxins) [35, 36]. The oxidative stress mediated by particulate matter (PM) may arise from: direct generation of reactive oxygen species (ROS) from the surface of soluble compounds; altered function of mitochondria or reduced nicotinamide adenine dinucleotide phosphate (NADPH)-oxidase; and activation of inflammatory cells capable of generating ROS and reactive nitrogen species (RNS), as well as oxidative DNA damage [37, 38]. The particle provides a template for electron transfer to molecular oxygen in these reduction and oxidation (redox) cycling events [39]. In addition, target cells, such as airway epithelial cells and macrophages, generate ROS in response to particle uptake by biologically catalysed oxidation reactions that occur in the cell membrane and mitochondria [4, 40–42]. *In vitro* studies have shown that inhaled PM causes expression of nuclear factor (NF)- κ B-related genes and oxidant-dependent NF- κ B activation [43, 44]. The dose of bio-available transition metal, rather than particulate mass, may be the primary determinant of acute inflammatory response [35, 37, 44]. However, other studies suggest that the hydrosoluble fraction is responsible for the oxidative damage to DNA [45]. The biological component of particles also seems to be related to oxidative stress [46], as well as bacterial endotoxin that induce the liberation of tumour necrosis factor (TNF)- α and interleukin (IL)-6 by macrophages [36].

Strong oxidative activity and the effective depletion of lung lining fluid antioxidants have been reported in large studies of ambient PM $<2.5 \mu\text{m}$ (PM_{2.5}) [17]. To defend against the oxidative damage, cells use up their stores of a key antioxidant, glutathione. The glutathione depletion can induce a state of cellular stress, which triggers an increase in the production of antioxidant enzymes through activation of a transcription factor nuclear factor-erythroid 2-related factor 2 [17]. Failure to overcome oxidative stress leads to the activation of additional intracellular signalling cascades that regulate the expression of cytokine and chemokine genes [15]. These products are produced locally in target tissues as well as systemically, and lead to widespread pro-inflammatory effects remote from the site of damage. In addition, PM appears to inhibit protective enzymes involved in oxidative stress responses depending on their toxicity (copper/zinc superoxide dismutase, manganese

superoxide dismutase, glutathione peroxidase and glutathione reductase) [47].

Diesel exhaust particles (DEPs) have a high content of elemental and organic carbon and are thought to be particularly toxic [15]. These particles consist of a carbon core with adsorbed organic compounds, such as polyaromatic hydrocarbons, quinones and redox-active metals, and the capacity of DEPs to induce oxidative stress is largely related to these adsorbed components. Animal experimental models, cell culture experiments and cell free systems involving DEPs have shown oxidative stress response and oxidative DNA damage. Human studies have shown increased neutrophils, B cells and alveolar macrophages in bronchoalveolar lavage fluid and an increased amount of pro-inflammatory cytokines, chemokines and adhesion molecules [48]. Exposure to DEPs has been shown to increase airway resistance, increase IL-6 and IL-8 in lavage fluid, increase IL-8 mRNA expression in bronchial mucosa and upregulate endothelial adhesion molecules P-selectin and vascular cell adhesion molecule-1 [49]. ROS formed at the epithelial level after DEP exposure upregulate IL-10, promoting antigen-presenting cells and allergy to pollen [15]. However, controlled exposure to DEP in human subjects has been shown to respond with an increase in low molecular antioxidants in the alveolar compartment [50]. The role of oxidative stress in response to DEPs and other particles is further supported by *in vitro* studies in which ROS are generated by macrophages, neutrophils, eosinophils and epithelial cells after stimulation by DEPs or particles [15]. Interestingly, low sulphur diesel combined with engine filters blocked a range of responses to DEPs including the oxidative stress responses in mice [51].

Alteration of autonomic functions also appears to be partly associated with oxidative stress [14]. Long-term exposure to low concentrations of PM_{2.5} has been shown to alter vasomotor tone, lead to vascular inflammation and potentiate atherosclerosis induced by high-fat chow in susceptible mice [52]. Although epidemiological evidence suggests that it is the fine (PM_{2.5}) or ultrafine (PM $<0.1 \mu\text{m}$) fraction that contains the toxic components; the large spectrum of disease end-points (from cardiovascular to asthma attack) suggest that more than one component may be driving the health effects [2].

O₃ is a very reactive gas whose uptake depends on the availability of antioxidants in the lining fluids, and its toxicity appears to be transmitted to the respiratory epithelium by secondary ROS formed by direct ozonisation of respiratory tract lining fluid lipids [16]. Alteration of the cell membrane translating an induction of lipid peroxidation and a significant modification of the redox status has been observed [53], as well as the activation of transcription factors such as NF- κ B and increased expression of a range of pro-inflammatory cytokines and adhesion genes [2, 6]. O₃ has been shown to react readily with ascorbic acid, uric acid and thiols, and exposure of these molecular species to O₃ results in their rapid depletion [6]. When these defence mechanisms are overwhelmed, O₃ may injure the underlying cells by inducing lipid peroxidation and activating inflammatory gene expression [6, 53]. Like O₃, nitrogen dioxide (NO₂) reacts with substrates present in the lung lining fluid compartment. The oxidised species arising from the reaction between NO₂ and lining fluid are responsible

for the signalling cascade of inflammatory cells into the lung [54–56].

A hierarchical oxidative stress model has been proposed to explain the dose-dependent response to air pollutant exposure [57]. Low exposure would lead to the formation of ROS activating an antioxidant response, followed by the transcription of enzymes important in detoxification, cytoprotective and antioxidant responses. These include phase II enzymes, whose induction serves as a detoxification mechanism (e.g. NAD(P)H:quinone oxidoreductase 1 (Nqo1) and glutathione S-transferase). At higher exposure, the transcription NF- κ B and activator protein-1 responses would be activated. This would lead to NF- κ B and mitogen-activated protein kinase signalling, altering the function of mitochondria or NADPH, and to increased expression of pro-inflammatory cytokines (such as TNF- α and IL-8 and IL-6) and genes coding adhesion molecules [2, 6, 15, 43, 44]. Any enhanced inflammatory response would lead to additional generation of ROS and RNS, together with oxidative DNA damage (fig. 1) [15, 37, 38].

Antioxidants and oxidative stress

Antioxidants in the lung are the first line of defence against oxygen free radicals. The respiratory tract lining fluids (RTLFL) contain a range of low molecular weight antioxidants similar to

those found in blood plasma, including reduced glutathione, ascorbic acid (vitamin C), uric acid and α -tocopherol (vitamin E). They also contain antioxidant enzymes, such as superoxide dismutase, glutathione peroxidase, thioredoxin reductase, catalase and the metal binding proteins ceruloplasmin and transferrin [2, 7]. All these antioxidants are free radical scavengers but also function as sacrificial targets for O₃ (ascorbate and urate) and react rapidly with this oxidant to limit its interaction with RTLFL lipids and proteins [58]. The composition and quantity of antioxidants in the RTLFL may represent an important determinant of individual responsiveness to air pollutants but should be thought of as a dynamic equilibrium with the antioxidant defences within the epithelium and the more remote plasma pool [59]. Controlled studies suggest that exposure to O₃ results in a depletion of RTLFL antioxidants followed by an enhancement of the movement of antioxidants to the RTLFL [60] or increased synthesis [3, 59]. Similarly, low-dose diesel exposure challenge in healthy volunteers was followed by an increase of inflammatory markers in bronchial lavage. No inflammatory response was seen in the alveolar compartment, but both reduced glutathione and urate concentrations were increased following diesel exposure suggesting differential antioxidant responses in the conducting airway and alveolar regions [50].

Although the inter-relation among antioxidant levels in RTLFL, cellular and plasma levels is not well understood, it appears that the susceptibility of the lung to oxidative injury depends largely on its ability to upregulate protective ROS- and RNS-scavenging systems and that the speed at which lost antioxidant defences can be replaced is a major determinant [58].

As many antioxidants are derived from the diet, several dietary factors have been implicated; mainly because of their potential role in inflammatory reactions. The following section will focus mostly on nutrients that have been used in supplementation studies to modulate the impact of air pollutants or might interact with the immune response. These factors include antioxidant vitamins, omega-3 fatty acids and other micronutrients that might affect the immune response.

Antioxidant nutrients

Vitamin C

Vitamin C, a water-soluble vitamin, is an abundant antioxidant substance and is widely distributed throughout the body including the extracellular lining fluid of the lung [17]. Ascorbate is an excellent reducing agent and scavenges free radical and oxidants. *In vitro* evidence suggests that vitamin C has a role as a chemical reducing agent both intracellularly and extracellularly. Intracellular vitamin C might prevent protein oxidation and regulate gene expression and mRNA translation. This is particularly relevant for the lung which is exposed to oxidative substances. Extracellular vitamin C protects against oxidants and oxidant-mediated damage [61]. It contributes to antioxidant activity through scavenging a variety of free radicals and oxidants, *in vitro*, including superoxide radical (O₂⁻), peroxy radicals, hydrogen peroxide, hypochlorous acid, singlet oxygen, oxidant air pollutants and oxidants that leak from activated neutrophils and macrophages [59, 61]. While the terminating product dehydroascorbate can be regenerated to ascorbate by intracellular enzymes, in particular thioredoxin




FIGURE 1. A model of the reaction of oxidants in the airway. Inhaled pollutants, such as ozone (O₃), nitrogen dioxide (NO₂), particulate matter <2.5 μ m (PM_{2.5}) or diesel exhaust particulates (DEPs), react with nonenzymatic antioxidant constituents of the respiratory tract lining fluid including: reduced glutathione (GSH); vitamin C; uric acid; and enzymatic antioxidants, such as extracellular superoxide dismutase (ecSOD), extracellular glutathione peroxidase (ecGSHpx) and thioredoxin. These molecules provide a protective screen against these pollutants. If defences are exceeded, the production of reactive oxygen species (ROS) is increased and oxidants may react with organic molecules, such as proteins or lipids, and alter the epithelium resulting in: cell activation and initiation of the inflammatory process; activation of neutrophils; and liberation of cytokines, chemokines and adhesion molecules. CHO: carbohydrate; NF- κ B: nuclear factor- κ B; AP-1: activator protein-1. Modified from [3].

reductase, which catalyses its regeneration [62], this regeneration is unlikely in the RTLf because of the lack of enzymes. Therefore, the maintenance of ascorbate level in the RTLf requires transportation from cellular sources or from the plasma pool [59]. Ascorbate also acts indirectly to prevent lipid peroxidation [59] and contributes to the regeneration of membrane-bound oxidised vitamin E [63]. Ascorbate plays a role in immune function and is transported into neutrophils and lymphocytes [18]. Whilst ascorbate has many antioxidant actions, it also has the capacity to act as a pro-oxidant in the presence of transition metals [64].

Vitamin E

Vitamin E, a lipid-soluble vitamin, represents the principal defence against oxidant-induced membrane injury in human tissue because of its role in breaking the lipid peroxidation chain reaction [64]. It is a potent peroxy radical scavenger and especially protects PUFAs within the phospholipid biological membrane and in plasma lipoproteins [65]. It also decreases production of prostaglandin E_2 , a metabolite of arachidonic acid produced by lipid peroxidation of lung cells after O_3 exposure [19]. Vitamin E appears to play a major role as an integral constituent of alveolar surfactant, whose quantity and composition conditions normal lung function [66].

β -Carotene

β -Carotene, a precursor to vitamin A and other carotenoids, accumulates in tissue membranes, scavenges $O_2^{\cdot-}$ and reacts directly with peroxy free radicals generated by O_3 [67]. It could, therefore, play a role in the control of inflammation and immune response through its antioxidant properties. However, recent research has shown that high-dose carotenoid supplementation may lead to both antioxidant and pro-oxidant reactions [68], depending on the redox potential of the biological environment in which it acts [69].

Other antioxidants, such as flavonoids, are scavengers of superoxide anions and peroxy radicals [70]. In addition to antioxidant activities, flavonoids can modulate cell signalling pathways [20]. Selenium, an essential trace element that plays a role in the detoxification of peroxides and free radicals [67], could also play an important role in the prevention of lung injury [21]. As an integral part of the glutathione peroxidases and thioredoxin reductase, selenium probably interacts with every nutrient that affects the pro-oxidant/antioxidant balance of the cell. It also appears to support the activity of vitamin E in limiting lipid oxidation [71].

Omega-3 PUFA

Increased intake of omega-3 PUFA (n-3 PUFA) can decrease the inflammatory reaction by changing the contents of lipid membranes and other substrates, which are in turn the substrates for eicosanoid production [72]. The substitution of n-3 PUFA (α -linoleic acid; 18:3n-3 and eicosapentaenoic acid (EPA); 20:5n-3) for n-6 fatty acids (linoleic acid; 18:2n-6) in the membrane leads to the production of less potent inflammatory mediators (prostaglandin E_3 instead of prostaglandin E_2 , and leukotriene 5 instead of leukotriene 4) [72]. Prostaglandin E_2 has been shown to act on T-lymphocytes to reduce the formation of interferon (IFN)- γ without affecting the formation of IL-4. This may lead to the development of allergic

sensitisation, since IL-4 promotes the synthesis of immunoglobulin E whereas IFN- γ has the opposite effect [73]. Leukotriene 4, a potent stimulator of airway smooth muscle cells, increases post-capillary vascular permeability and mediates asthma by vasoconstriction and mucus secretion. The competitive interactions between n-6 PUFA and n-3 PUFA determine the cellular contents of arachidonic acid and EPA.

Increased intake of n-3 PUFA appears to decrease the risk of sudden and nonsudden death from myocardial infarction and nonfatal myocardial infarction [74–76]. The protective effect of n-3 PUFA may be linked, in part, to its cardiac and arrhythmic effects, including increasing heart rate variability (HRV) [22, 74, 77]. There is a positive correlation between the baseline cell membrane concentrations of n-3 PUFA and the degree of HRV, both in healthy subjects and in patients with coronary artery disease [23, 78]. Along with increasing HRV, other anti-arrhythmic mechanisms of n-3 PUFA have also been described, including the capacity to stabilise the electrical activity of cardiac myocytes by modulating sarcolemmal ion channels and voltage-dependent sodium channels [24], and the capacity to reduce myocardial infarct size in animal models of ischaemia and reperfusion [24]. N-3 PUFA also appear to: decrease the risk of thrombosis; decrease serum triglyceride levels, slowing the growth of atherosclerotic plaque; improve vascular endothelial function; lower blood pressure; and decrease inflammation [79].

Other micronutrients and immune functions

Micronutrients such as zinc, vitamin A and folic acid can also influence several components of immunity, altering the function of macrophages and thus their role in innate immunity and inflammation. Studies have shown that deficiencies in these micronutrients can significantly alter macrophage phagocytosis and their production of cytokines (IL-1 and IL-6, TNF- α and IFN- γ). These deficiencies also alter natural killer cell function, neutrophil motility and antimicrobial activity [25].

Nutrient supplementation and effects of air pollution

The effects on air pollutant toxicity of nutrient supplementation at levels higher than is physiologically required have been studied in both animals and humans and summarised previously [2, 11, 17, 80].

Experimental animal studies

Results of animal studies suggest that supplementation with vitamin C and vitamin E modulates the pulmonary response to exposure to photo-oxidants, such as O_3 or NO_2 [17, 81], and that vitamin C, uric acid and glutathione located in the respiratory tract lining fluid are consumed on exposure to O_3 and NO_2 [16, 82, 83]. Dietary deficiency of vitamin C appears to quickly translate to decreased levels of vitamin C in blood and RTLf [84]. Temporary vitamin E deficiency may induce reversible changes in the expression of pro-inflammatory markers, reduce surfactant lipid synthesis in alveolar type II cells and favour the development of injury in response to air pollution insults [66]. Further experimental studies using antioxidants, iron chelators or other substances support the role of ROS as mediators of the effects of particulates [37, 54]. Oxidative stress appears to play a critical role in the activation

of NF- κ B, and cytokine-induced NF- κ B activation is prevented after treatment with antioxidants or metal chelators [54]. N-acetylcysteine, a powerful antioxidant, had a protective effect on inflammatory response and oxidative stress damage in rats exposed to coal dust [85] and on changes in heart rate and decrease in HRV in rats exposed to urban air particles [86].

Human studies

There is little information on the impact of antioxidant supplementation on the acute effects of air pollution exposure in humans. Most existing studies have focused on the changes of acute lung function. Other outcomes included bronchial airway reactivity, inflammatory response and changes in HRV but are less numerous and consistent. All these studies were experimental studies using supplements.

Antioxidant supplementation

Lung function and airway reactivity

Early studies used experimental protocols with single pollutants and a small number of healthy adults. Levels of O₃ and NO₂ were very high (usually close to 1,000 $\mu\text{g}\cdot\text{m}^{-3}$ and >3,000 $\mu\text{g}\cdot\text{m}^{-3}$, respectively) and subjects were supplemented for a relatively short period of time with high doses of vitamin C or vitamin E (eight to 16 times the USA recommended daily allowance of vitamin C (60 $\text{mg}\cdot\text{day}^{-1}$) and vitamin E (8 $\text{mg}\cdot\text{day}^{-1}$)) [2, 87–89]. A modulating effect of antioxidant supplementation was observed in some studies of acute lung function changes [89] and airway reactivity [87] but not in others.

More recent experimental studies have addressed conditions in which the O₃ level and supplement doses were lower. In a study of asthmatic adults, a cocktail of vitamin C (500 mg) and vitamin E (400 UI) protected against a decrease in peak expiratory flow from SO₂ challenge after O₃ exposure [90]. In another study [91], subjects were first deprived of vitamin C and then supplemented with a relatively low dose of vitamin C (250 mg), vitamin E (100 mg) and vegetable cocktail. Supplementation protected against acute change in lung function (forced expiratory volume in one second and forced vital capacity) after O₃ challenge. However, in well nourished individuals sensible to O₃, supplementation with vitamin C (500 mg) and vitamin E (100 mg) provide no protective effect on inflammatory response or lung function decrease after O₃ challenge. This lack of protection was observed despite elevated plasma vitamin C (+60.1%) and vitamin E (+51.4%) concentrations following supplementation, and increased vitamin C concentrations in the airways after supplementation following O₃ exposure [92].

Supplementation studies conducted in free-living populations of healthy exercising adults (the Netherlands) or adults exposed to high levels of air pollutants (Mexico) support the hypothesis that antioxidant supplementation protects against the acute effects of O₃ on lung function. In these studies, healthy adults were randomised to receive vitamin C (650 mg), vitamin E (75 mg) and β -carotene (15 mg) for several weeks [80, 93–95]. More recently, a study of asthmatic children exposed to high levels of air pollutants in Mexico City also suggested that supplementation with vitamin C (250 $\text{mg}\cdot\text{day}^{-1}$) and vitamin E (50 $\text{mg}\cdot\text{day}^{-1}$) had a modulating effect on acute lung function changes [96]. The positive effect of antioxidant

supplementation was mostly found in children genetically susceptible to the effects of oxidants (glutathione S-transferases (GST)M1 null genotypes) [97].

Inflammatory response

Only three studies have evaluated the impact of antioxidant supplementation on airway inflammatory response to air pollutant exposure. SAMET *et al.* [91] observed no difference in the bronchoalveolar lavage content of polynuclear cells and other inflammatory markers between supplement and placebo groups after O₃ challenge. Similarly, Mudway *et al.* [92] reported no effect of supplementation with vitamin C and vitamin E on O₃-induced neutrophilia in healthy individuals responsive to O₃. In contrast, asthmatic children heavily exposed to air pollutants and supplemented with vitamin C and vitamin E had significantly lower levels of IL-6 and IL-8 in nasal lavage than children receiving placebo [98].

n-3 PUFA supplementation

Lung function and inflammatory response

The impact of n-3 PUFA supplementation on asthmatic symptoms and exercise-induced bronchoconstriction has been examined among asthmatic subjects in various recently reviewed studies [12, 34, 99]. Most of these studies enrolled a small number of asthmatic patients randomly assigned to receive a high dose of n-3 PUFA (3–4 g of EPA) for a short time-period (6–10 weeks); results were inconsistent. Studies with longer intervention periods, from 6 months to 1 yr, also led to inconsistent results with some studies showing improvement in lung function [100, 101] or inflammatory markers [101–103], or no effect [104]. The dosage and duration of n-3 PUFA supplementation, and the type of asthmatic patients differed between studies and may explain the discrepancy between these studies [12, 34]. The Cochrane database of systematic reviews identified 22 studies but included only nine that fulfilled the inclusion criteria and concluded that data were insufficient to determine the effect of n-3 PUFA in asthma. None of these studies include information on air pollution.

Cardiovascular effect

Increased intake of n-3 PUFA either from dietary sources or as a pharmacological supplementation has been shown to decrease the risk of mortality from coronary heart disease [105]. In a randomised trial conducted in nursing home residents, supplementation with 2 $\text{g}\cdot\text{day}^{-1}$ of fish oil (each 1 g capsule contained 83.2 % of omega-3 fatty acids) significantly decreased the effect of PM_{2.5} on time and frequency domain parameters of HRV [106]. This is one of two studies providing evidence that oxidant stress is one of the mechanisms explaining the effect of particle air pollution on the cardiovascular system [107]. The other study reported that statins had a mitigating effect on the HRV effects of particulate air pollution in subjects genetically susceptible to oxidative stress (lacking the GSTM1 allele) [108].

Modifiers of an individual's response to oxidative stress

Under the biological model of oxidative stress one would expect factors that modify the response to oxidative stress to also alter the effects of air pollution. Thus, nutritional status, chronic diseases and genetic factors are candidates to

determine susceptibility to oxidative stress-related effects of air pollution [26] as all these conditions are related to poor antioxidant defence.

Nutritional status

Antioxidant vitamin supplementation provides some protection against the adverse effect of O₃ on lung function in asthmatic children with slight deficiencies in these nutrients [96], and to adults depleted in vitamin C [91]. In contrast, vitamin supplementation did not protect against O₃-induced lung function decrement in well nourished subjects [109].

Chronic diseases

Most chronic diseases are associated with chronic inflammation [13, 27, 28, 110–112], which might increase susceptibility to the additional oxidative stress caused by air pollution exposure. In particular, subjects with asthma [29], chronic obstructive lung diseases [113], diabetes [114] and cardiovascular diseases [115] have all been shown to have antioxidant deficiency [13] and be more susceptible to the effects of air pollution [108, 115]. As observed in the case of cigarette smoke, a significant source of oxidative stress, air pollutants would lower antioxidant defences, with deleterious health consequences [116, 117]. Evidence of the potential beneficial effect of antioxidants can be found in studies of elderly subjects in which treatment with statins [108] and n-3 PUFA supplementation [106] had a beneficial effect on response to particulate exposure.

Genetic susceptibility

As oxidative stress is an important pathway activated/involved in the adverse effects of air pollution, the genes involved are of primary interest. Most studies have focused on single gene polymorphisms; however, it is likely that there will be a hierarchy of genes determining susceptibility, rather than one individual gene driving this process [15].

GST enzymes: GSTM1, GSTP1

GST are phase II xenobiotic metabolising enzymes that participate in the detoxification of ROS by catalysing their conjugation with glutathione [118, 119]. The common null allele of *GSTM1* results in a complete lack of the enzyme and reduced or no conjugation activity [120]. It has been associated with an increase in asthma and wheezing among children exposed to environmental tobacco smoke *in utero*, with a decrease in lung function growth [121, 122], and also with a rapid decline in lung function in smokers [123]. In addition, polymorphic *GSTM1* has been shown to act as a modifier of the lung response to fire smoke [124] and O₃ [125]. Antioxidant supplementation with vitamin C and E appears to modulate the effect of O₃ in asthmatic children homozygous for the *GSTM1* null allele [97]. Allergen sensitive subjects with low responsive genotypes show enhanced susceptibility to the adjuvant effects of DEP [126]. A *GSTM1* polymorphism has also been shown to increase sensitivity to PM, as evidenced by greater changes in HRV [108]. Moreover, glutaryl coenzyme A inhibitors, *i.e.* statins, with known antioxidant and anti-inflammatory properties mitigate against the effects of ambient particles on HRV in subjects lacking the *GSTM1* allele [107, 108].

Other genes

The Toll-like receptor 4 (*TLR4*; *xr 4*) gene has been implicated in innate immunity and endotoxin susceptibility [127] and has been hypothesised to play a role in O₃-induced hyperpermeability [26]. TNF- α (*Xr17*) has been related to lung function changes after O₃ exposure [128] and to an increased risk of asthma and wheezing that can be modified by O₃ exposure [129]. TNF has been identified as a candidate gene for O₃-induced airway inflammation and hyperresponsiveness [130]. Polymorphisms in TNF and lipoteichoic acid have been associated with respiratory effects of O₃ in humans [128]. *Arginase II* has been associated with an increased risk of asthma in children, and the association appeared stronger among children with a smoking parent [131] suggesting that air pollutants could also play a role.

Gene-gene interactions

O₃-induced acute effects on respiratory function have been shown to be smaller in subjects with *GSTM1* null and *NOQ1* Pro/Pro genotypes [132]. Similarly, a study examining asthma risk in a population highly exposed to O₃ showed that the risk of asthma was significantly associated with the *NOQ1* genotype in subjects with the null genotype for *GSTM1* [133]. Both genes have a specific function in antioxidative activities.

FURTHER EPIDEMIOLOGICAL RESEARCH

There is now substantial evidence that air pollution exposure results in increased oxidative stress, alterations in immune regulation and repeated inflammatory responses that overcome lung defences to disrupt the normal regulatory and repair processes [10, 15]. As summarised previously, despite a plausible mechanistic model linking air pollution, oxidative stress and dietary supplementation, evidence is not sufficient. Further randomised controlled trials (RCTs) are needed in order to better understand the potential protective effect of nutrient supplementation on the effect of air pollution on respiratory and cardiovascular functions and inflammatory responses.

RCTs provide a good alternative to maximise contrast in nutrient intake for evaluating the interaction of dietary factors and air pollutants and should be conducted in both the controlled setting and in free-living populations. A controlled setting will allow assignment of air pollutant exposure and, therefore, provide an accurate representation of the health effects and potential modulating effects of supplementation, while RCT conducted in free-living populations will have the advantage of representing real-life conditions.

Susceptible subjects, such as those with pre-existing respiratory or cardiac disease, micronutrient deficiency or genetic susceptibility, are the most likely to benefit from nutritional intervention (see Modifier of response section); therefore, RCTs should focus on these population subgroups. Short- and long-term effects can be studied; however, the major challenge in long-term effect studies is to assess the appropriate time-frame of exposure for the induction of the disease and, therefore, the relevant period and duration of the supplementation. There is accumulating evidence that exposure during lung development in foetal life and early childhood plays a major role, as in the case of maternal smoking [134–136]. Therefore, RCTs of pregnant females with specific risks (such as asthmatic or

TABLE 1 Biomarkers of oxidative stress most commonly used in clinical and epidemiological studies

Type of measurement	Biomarker	Biological sample	Laboratory technique	Sensitivity and specificity	Comments	[Ref.]
TAC	TRAP	Plasma	Fluorescence	Good	Measures the cumulative action of all antioxidants present in plasma and body fluids TRAP: indirect measure TRAP+R-PE: direct measure of peroxy radical attack on R-PE. Affected by protein concentration Plasma better than serum	[140–142]
	TRAP + R-PE	Serum		Possible artefactual confounding		
Lipid peroxidation	TBARS	Tissue Plasma Serum	Spectrophotometry Colourimetry Fluorometry	Low specificity	Easy to use Indirect measure	[143, 144]
	MDA-TBA derivatisation	Plasma	TBARS	Low specificity	Measures MDA, end product of lipoperoxidation. MDA is generated mainly by arachidonic acid and docosahexaenoic acid With HPLC detection, MDA is not a specific product of lipid peroxidation	[143–145]
		Serum	HPLC/MS	Good		
	Free MDA	Plasma Serum	HPLC HPCE	Good	Low amount of plasma needed Fast and practical for clinical measurements Low detection limit	[145, 146]
				Good		
	4-hydroxynonenal 4-hydroxy	Tissue	ELISA	Good	HNE is a toxic product of lipid peroxidation and second toxic messenger of free radicals	[147, 148]
		Blood Urine	GC/MS			
	Hydrocarbons: ethane and pentane	EBC	GC	Pentane: low specificity	Hydrocarbons are produced through peroxidation of fatty acids in cellular biomembranes, by ROS Ethane: faster chromatographic measurement compared with other hydrocarbons; better marker for lipid peroxidation Background level of pentane and isoprene in human breath difficult to separate pentane from isoprene by chromatography Possible contamination with ambient air ethane and pentane	[149–152]
Ethane: good						
Conjugated dienes	Plasma Serum	Spectrophotometry HPLC	Validity still questionable	Other biological substances, even polyunsaturated fatty acids, absorb in the same UV region CD generation continues <i>ex vivo</i> after sampling Plasma CD is >90% derived from 9, 11 diene-conjugated linoleic acid from dietary dairy products	[150, 153]	
LDL oxidation	Plasma	<i>Ex vivo</i> LDL by CD assay with spectrophotometric determination	Good	Measures the rate of CD formation Cannot be known for certain whether the <i>in vitro</i> situation accurately reflects <i>in vivo</i> events Should reflect the antioxidant defence system. Vitamin E has shown reasonably consistent effects in increasing the resistance of LDL to oxidation	[143, 153–155]	
			Good			
	Plasma Serum	<i>In vivo</i> LDL-BDC with spectrophotometric determination	Good	Faster and simpler to perform than the <i>ex vivo</i> procedure Measures amount baseline diene conjugation	[156]	
Oxidised LDL	Plasma	ELISA	Poor	These modifications may occur independently of lipid peroxidation Still unclear whether it can serve as a peripheral marker High variability	[144, 152, 157]	
Lipid hydroperoxides: CEOOH	Plasma	HPLC assay with chemiluminescence detection	Not confirmed	Not detectable in young healthy controls Direct indicator of lipid peroxidation	[144, 158]	

TABLE 1 Continued.

Type of measurement	Biomarker	Biological sample	Laboratory technique	Sensitivity and specificity	Comments	[Ref.]
Eicosanoids	F2-isoprostane	Plasma	HPLC	Good	These markers reflect respiratory tract integrity between reactive nitrogen species and ROS Interaction with other prostanoids Potent biological activity 8-iso-PGF _{2α} is a major component of total F ₂ isoprostanes In plasma, possibility of artefactual generation due to arachidonic acid autoxidation Better in urine - less interaction	[143, 144, 152, 159, 160]
		Serum	GC/MS			
		Urine	ELISA			
		EBC				
	PGE ₂	EBC	HPLC/MS/MS	Good	Not flow dependent in healthy subjects	[159–162]
		Plasma Sputum	ELISA GC/MS			
LTB ₄	EBC Plasma Serum Urine Sputum BAL	EBC	GC/MS	Good	LTB ₄ is a potent neutrophil chemoattractant and may contribute to airway narrowing by producing local oedema and increasing mucus secretion	[159–161]
		Plasma	HPLC			
		Serum	ELISA			
		Urine				
		Sputum BAL				
Nitrogen reactive species	Nitrite: NO ₂ ⁻ Nitrate: NO ₃ ⁻	EBC	Colourimetry	Good	In healthy children, nitrite values are not related to levels of exhaled NO Both nitrite and nitrate quantification	[159, 163–166]
		Plasma	Fluorometry Ionic chromatography GC/MS HPLC			
	S-nitrosothiols 3-nitrotyrosine	Plasma BAL	Fluorometry GC/MS			
DNA oxidation	8-OHdG	Urine	ELISA	Poor	May be influenced by the metabolic rate and also by excision repair GC/MS: level of 8-OHdG overestimated ELISA values higher than HPLC values	[143, 170–173]
		DNA	CG/MS HPLC/ECD			
	8-oxoGua	DNA	CG-MS	Good	HPLC-ECD generally yields lower values Enzymatic approach: FPG may detect lesions other than 8-oxo-7, 8-dihydroguanine; the method relies on indirect calibration Reported strong correlation between overnight and 24 h urinary 8-oxodGuo [#]	[174, 175]
			HPLC-ECD			
			HPLC-MS Comet assay ELISA			
	8-oxodGuo	24 h urine	CG-MS	Good	HPLC-ECD generally yields lower values Enzymatic approach: FPG may detect lesions other than 8-oxo-7, 8-dihydroguanine; the method relies on indirect calibration Reported strong correlation between overnight and 24 h urinary 8-oxodGuo [#]	[174, 175]
			HPLC-ECD HPLC-MS Comet assay ELISA			
Modified comet assay	DNA	SCGE	Good	Measures DNA strand breaks Proportion of DNA in the tail indicates the frequency of breaks Particularly sensitive to oxidative attack by H ₂ O ₂	[143, 176]	
HmdU	Plasma Serum	ELISA	Good	Autoantibody to oxidised DNA Product of thymine oxidation	[143, 177, 178]	
Protein oxidation	Protein carbonyl	Plasma Lung aspirate	Colourimetric method ELISA HPLC	Good	Measures generic oxidation; does not differentiate between those protein carbonyl arising directly from protein oxidation and those formed by adduction of other oxidised products	[143, 153, 179]

TABLE 1 Continued.

Type of measurement	Biomarker	Biological sample	Laboratory technique	Sensitivity and specificity	Comments	[Ref.]	
Other	GSH	Sputum	Spectrophotometry	Good	GSH is a protective antioxidant against oxidative stress Level of GSH depends on biological sample	[159, 180–184]	
		Plasma Saliva					
	GSH/GSSG ratio	Plasma Serum	BAL	Reverse phase HPLC	Good	Decrease in GSH/GSSG indicates chronic oxidative stress	[153, 185]
			EBC	HPLC /with fluorescence detection	Good		
	H ₂ O ₂	EBC		Spectrophotometry	Poor: high variation	Concentration appears to be expiratory flow rate dependent Wide variability in mean exhaled H ₂ O ₂ concentration in healthy nonsmoking adults Other factors: exercise, food, beverage intake	[159, 186–188]
			Fluorometry Chemiluminescence				
CC16	Serum BALF		Latex immunoassay	Good	These tests evaluate the integrity of respiratory tract Peripheral marker CC16 protects the respiratory tract against oxidative stress and inflammation	[189–192]	
			ELISA				
Thioredoxin	Serum		ELISA	Good	Thioredoxin is induced by oxidative stress and secreted by cells	[193–195]	

TAC: total antioxidant capacity; TRAP: total radical trapping antioxidant parameter; R-PE: R-phycoerythrin; TBARS: thiobarbituric acid-reactive substances; MDA-TBA: malondialdehyde-thiobarbituric acid; HPLC: high performance liquid chromatography; MS: mass spectrometry; EBC: exhaled breath condensate; UV/Vis: UV/visible detection; HPCE: high performance capillary electrophoresis; HNE: 4-hydroxynonenal; GC/MS: gas chromatography/MS tandem; ROS: reactive oxygen species; CD: conjugated dienes; LDL: low-density lipoprotein; BDC: baseline diene conjugation; CEOOH: cholesteryl ester hydroperoxides; PG: prostaglandin; LTB₄: leukotriene B₄; BAL: bronchoalveolar lavage; NO: nitric oxide; 8-OHdG: 8-hydroxy-2'-deoxyguanosine; ECD: electrochemical detection; 8-oxoGua: 8-oxo-7,8-dihydroguanine; FPG: fasting plasma glucose; SCGE: single cell microgel electrophoresis; 8-oxodGuo: 8-oxo-7,8-dihydro-2'-deoxyguanosine; HmdU: 5-hydroxymethyl-2'-deoxyuridine; GSH: reduced glutathione; GSSG: oxidised glutathione (disulfide form); NL: nasal lavage; BALF: BAL fluid(s). #: r=0.93, p<0.01.

atopic mothers) might provide some insight into the role of antioxidants and n-3 PUFA as modulators of the air pollution effect. In these studies, a major challenge is the accurate assessment of air pollution exposure, oxidative stress, biomarkers of nutritional status and health outcomes. Standardisation of these factors within and between studies is crucial to allow comparability of results. In the following section some issues to be considered in future studies will be discussed.

Air pollution exposure

Contrasts in exposure need to be maximised to be able to distinguish between effects in the placebo group and smaller or no effects in the supplemented groups. Depending on the study design and hypotheses tested, either temporal or spatial contrast should be large. Multicentre studies including areas with contrasting air pollution levels and the enrolment of random samples of participants within each centre might be an option. Moreover, the design of the exposure assessment must take into account the relationship between measured or measurable markers of oxidant pollution and personal exposure to the pollutant relevant to the hypothesis. For example, there are often large indoor/outdoor ratios in O₃ concentrations and these can be very heterogeneous across homes. Personal O₃ concentration may be very poorly correlated with ambient

levels in certain areas. It might be useful to measure the redox activity of ambient pollutants or the antioxidant depletion rates, as these may be the most relevant characteristics in the hypothesised pathways of redox imbalance. Various assays have been developed to measure the redox activity of particles, such as OH radical formation or antioxidant depletion rates [137]. However, the measurement methods may need further development to be applicable in epidemiological studies, in particular, for personal exposure assessment.

Biomarkers of oxidative stress

The advantage of using biomarkers is that they integrate both the effects of oxidant exposure and the full range of antioxidant protective mechanisms *in vivo* [30]. However, samples can be oxidised during handling, processing and analysis, so there is potential for artefacts in estimates of baseline levels of oxidation markers. The magnitude of this problem varies between biomarkers [31, 138]. Most of these biomarkers include measures of lipid, DNA and protein oxidation. Recent review articles provide broad coverage of this topic [30, 139]. Table 1 presents a summary of oxidative stress biomarkers useful for clinical and epidemiological studies including: the type of marker; the biological media for measurement; the laboratory techniques most frequently used; an appreciation of its

TABLE 2 Biomarkers of nutrient intake most commonly used in clinical and epidemiological studies

Type of measurement	Biological sample	Laboratory technique	Comments	Characteristics and sources	[Ref.]
Carotenoids	Serum	HPLC	Poor bioavailability in raw food, improved by mild cooking or heating (e.g. lycopene in tomato juice)	Liposoluble	[143, 196–198]
β -Carotene	Plasma		Reflect short-term intake	Red, orange and yellow fruits and vegetables (sweet potato, carrots, winter squash)	
α -Carotene	Induced sputum		Need to control for cholesterol level	Green vegetables	
Lycopene	Adipose tissue		Adipose tissue reflects long-term exposure		
Lutein			May not reflect concentration in target tissue		
Xanthine					
β -Cryptoxanthin					
Tocopherols	Serum	HPLC	Serum and plasma reflect short-term intake	Liposoluble	
α -Tocopherol	Plasma		Need to control for cholesterol level	Vegetable and seed oils (corn, safflower, soy)	[143, 199]
γ -Tocopherol	Adipose tissue		Adipose tissue reflects long-term exposure	Beans, eggs, green vegetables	
Vitamin C	Serum	HPLC	Vitamin C in food can be destroyed by exposure to high temperature, oxidation or cooking in large amount of water	Hydrosoluble	
	Plasma		Response to intake up to 50–90 mg·day ⁻¹ , then eliminated by renal clearance	Fruits: papaya, cantaloupe, citrus fruits, strawberries	[143, 200]
			Reflects short-term intake	Vegetables: cauliflower, broccoli, brussel sprouts, kale, sweet peppers	
			Predicts intake at low level of vitamin intake		
Selenium	Plasma	Atomic absorption	At higher levels of intake, the correlation between plasma selenium concentration and dietary intake depends on the chemical form of selenium in the diet	Cereals and grains	[143, 201–204]
	Toenail	spectrophotometry		Animal products, especially organ meats and seafood	
		HPLC	Selenium content of cereals and grains depends on the soil content		
			Plasma reflects short-term intake		
			Nail and whole blood reflect long-term exposure (>26–56 weeks)		
Flavonoids	Serum	HPLC	Measures the usual dietary intake over 1 week	Apples, lemons, oranges	[205, 206]
	Urine			Potatoes, cauliflower	
				Tea	
				Skin of tubers and roots	
				Red wine	
Isoflavonoids	Serum	GC/MS	Sex differences in metabolism and excretion	Legumes: soybeans, beans, lentils, chickpeas.	[207–209]
	Urine	HPLC			

TABLE 2 Continued.

Type of measurement	Biological sample	Laboratory technique	Comments	Characteristics and sources	[Ref.]
Lignans	Serum -24–72 h urine	HPLC	Sex differences in metabolism and excretion	Oil seeds (flax seed, soybean, rapeseed) Whole-grain cereals (wheat, oats, rye), legumes, vegetables; fruits	[207–209]
PUFA	Free fatty acids in serum or plasma	HPLC	Samples are temperature and oxygen sensitive	Fish oils	[210–212]
n-3 PUFA	Components of circulating triglycerides	GC/MS	Potential for oxidation and degeneration over time	Fish and shellfish	
n-6 PUFA	Components of circulating phospholipids	GLC	Free fatty acids, phospholipids and cholesterol esters represent the intake over the last few days or meals	Soy and canola oil	
	Cholesterol esters		Serum fatty acids appear to be sensitive to changes in diet; high fluctuation (10–12%) and lab error <5%		
	Red blood cell membranes		Components of triglycerides represent intake over the past few hours		
	EBC		RBC reflect longer term intake (half-life of RBC: 120 days)		
	Adipose tissue		RBC sample: collected whole blood is suspended in phosphate buffer and centrifuged; packed red cells are stored at -80°C		
			RBC: may contain lower levels of n-3 and n-6 PUFA		
			Adipose tissue reflects long-term intake if no severe weight loss has occurred		
Folate	Serum	ELISA	Serum: short-term folate	Leafy vegetables	[213, 214]
	RBC		RBC: dietary intake over last 120 days	Dry beans and peas, fortified cereal Some fruits	
Zinc	Plasma	Atomic absorption	Plasma: most frequently used	Oysters	[202, 215–218]
	Cells	spectrometry	Possibility of no association between zinc intake and plasma zinc	Animal proteins	
	Erythrocyte, monocyte, neutrophil, platelet		Cells: complex sample preparation	Beans	
	Hair		Poor sensitivity, imperfect specificity	Nuts	
	Nails			Pumpkin and sunflower seeds	
	Urine				

HPLC: high performance liquid chromatography; GC: gas chromatography; MS: mass spectrometry; PUFA: polyunsaturated fatty acids; GLC: gas liquid chromatography; EBC: exhaled breath condensate; RBC: red blood cells.

sensitivity and specificity based on the literature review; and some additional comments [140–195].

Biomarkers of exposure to antioxidant nutrients and n-3PUFA

These biochemical indicators have the advantage of integrating different food sources and providing a better estimation of the internal dose, *i.e.* a closer indication of the amount of nutrient available after absorption and metabolism [33]. They can also be used in intervention studies to monitor compliance with the supplement. However, they are subject to measurement errors and sampling, storage, handling and laboratory analysis and temporality issues need to be carefully considered [30]. Table 2 presents a summary of biomarkers of antioxidant and n-3 PUFA intake used in clinical and epidemiological studies including: the type of marker; the biological media for measurements; the laboratory techniques most frequently used; the characteristics and food sources of these nutrient biomarkers; and some additional comments [196–218].

Health end-points

The limited validity of symptoms of respiratory or cardiac diseases has been extensively discussed [219, 220]. Objective outcomes, such as lung function, nitric oxide in exhaled breath, carotid intima-media thickness, electrocardiographic abnormalities or HRV, are less prone to bias and may be a good alternative but their long-term predictive value is uncertain. Biological indicators, such as pro-inflammatory markers (*e.g.* IL-6, IL-4, TNF- α , IFN- γ) in sera, exhaled breath and nasal lavage, and peripheral inflammatory markers (*e.g.* cell counts, fibrinogen, C-reactive protein, von-Willebrand factor, prostaglandin E2, plasminogen activator inhibitor, cell adhesion molecules) might provide useful information about potential mechanisms of air pollutant exposure. However, they are subject to large within-person variability and limited specificity as they are common to different end-points; therefore, serial measurements over the study period are required. In addition, intra-individual variability and the temporal frame need to be considered for any of the transient end-points. A mechanistic approach that includes evaluation of several end-points at the clinical and biological levels seems most appropriate. Further understanding of the crucial role of transcription factors, DNA methylation and RNA control of gene expression will provide new perspectives on the complex interaction of air pollutants and nutritional factors.

CONCLUSION

Oxidative stress is one of the main mechanisms by which air pollutants affect respiratory and cardiovascular health. Short-term randomised supplementation trials suggest that antioxidant vitamins and n-3 polyunsaturated fatty acids might protect against the acute effect of these pollutants, particularly in vulnerable subgroups [80, 96, 106]. However, the evidence is still limited because of the small sample size in most studies and the lack of comprehensive assessment of baseline nutritional status and oxidative stress response. Future studies should include randomised control trials of antioxidant or n-3 polyunsaturated fatty acid supplementation in susceptible populations and measure clinical, as well as intermediate, outcomes and biomarkers of oxidative stress and nutrient

intake considering factors, such as reproducibility, inter- versus intra-person variability, detection limits and specificity and sensitivity of these markers. Doses and duration are still under debate but harmonisation between studies is desirable for comparison purposes.

ACKNOWLEDGEMENTS

The authors would like to thank G. Evans for reviewing the English of the manuscript.

REFERENCES

- 1 Brunekreef B, Holgate ST. Air pollution and health. *Lancet* 2002; 360: 1233–1242.
- 2 Kelly FJ. Oxidative stress: its role in air pollution and adverse health effects. *Occup Environ Med* 2003; 60: 612–616.
- 3 Mudway IS, Kelly FJ. Ozone and the lung: a sensitive issue. *Mol Aspects Med* 2000; 21: 1–48.
- 4 Nel A. Atmosphere. Air pollution-related illness: effects of particles. *Science* 2005; 308: 804–806.
- 5 Schlesinger RB, Kunzli N, Hidy GM, Gotschi T, Jerrett M. The health relevance of ambient particulate matter characteristics: coherence of toxicological and epidemiological inferences. *Inhal Toxicol* 2006; 18: 95–125.
- 6 Cross CE, Valacchi G, Schock B, *et al.* Environmental oxidant pollutant effects on biologic systems: a focus on micronutrient antioxidant-oxidant interactions. *Am J Respir Crit Care Med* 2002; 166: Suppl. 12, S44–S50.
- 7 Bowler RP, Crapo JD. Oxidative stress in allergic respiratory diseases. *J Allergy Clin Immunol* 2002; 110: 349–356.
- 8 Misso NL, Thompson PJ. Oxidative stress and antioxidant deficiencies in asthma: potential modification by diet. *Redox Rep* 2005; 10: 247–255.
- 9 Pryor WA. Mechanisms of radical formation from reactions of ozone with target molecules in the lung. *Free Radic Biol Med* 1994; 17: 451–465.
- 10 Gilliland FD, McConnell R, Peters J, Gong H Jr. A theoretical basis for investigating ambient air pollution and children's respiratory health. *Environ Health Perspect* 1999; 107: 403–407.
- 11 Pryor WA. Can vitamin E protect humans against the pathological effects of ozone in smog? *Am J Clin Nutr* 1991; 53: 702–722.
- 12 Woods RK, Thien FC, Abramson MJ. Dietary marine fatty acids (fish oil) for asthma in adults and children. *Cochrane Database Syst Rev* 2003; 2: CD001283.
- 13 Boots AW, Haenen GRMM, Bast A. Oxidant metabolism in chronic obstructive pulmonary disease. *Eur Respir J* 2003; 22: Suppl. 46, 14S–27S.
- 14 Brook RD, Brook JR, Rajagopalan S. Air pollution: the "Heart" of the problem. *Curr Hypertens Rep* 2003; 5: 32–39.
- 15 Saxon A, Diaz-Sanchez D. Air pollution and allergy: you are what you breathe. *Nat Immunol* 2005; 6: 223–226.
- 16 Pryor WA, Squadrito GL, Friedman M. A new mechanism for the toxicity of ozone. *Toxicol Lett* 1995; 82–83: 287–293.
- 17 Hatch GE. Asthma, inhaled oxidants, and dietary antioxidants. *Am J Clin Nutr* 1995; 61: Suppl. 3, 625S–630S.

- 18 Bast A, Haenen GR, Doelman CJ. Oxidants and antioxidants: state of the art. *Am J Med* 1991; 91: Suppl. 30, S2–13S.
- 19 Singh U, Devaraj S, Jialal I. Vitamin E, oxidative stress, and inflammation. *Annu Rev Nutr* 2005; 25: 151–174.
- 20 Williams RJ, Spencer JP, Rice-Evans C. Flavonoids: antioxidants or signalling molecules? *Free Radic Biol Med* 2004; 36: 838–849.
- 21 Romieu I. Nutrition and lung health. *Int J Tuberc Lung Dis* 2005; 9: 362–374.
- 22 Routledge HC, Ayres JG, Townend JN. Why cardiologists should be interested in air pollution. *Heart* 2003; 89: 1383–1388.
- 23 Leaf A, Kang JX, Xiao YF, Billman GE. Clinical prevention of sudden cardiac death by n-3 polyunsaturated fatty acids and mechanism of prevention of arrhythmias by n-3 fish oils. *Circulation* 2003; 107: 2646–2652.
- 24 Kris-Etherton PM, Harris WS, Appel LJ, American Heart Association Nutrition Committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation* 2002; 106: 2747–2757.
- 25 Erickson KL, Medina EA, Hubbard NE. Micronutrients and innate immunity. *J Infect Dis* 2000; 182: Suppl. 1, S5–S10.
- 26 Kleeberger SR. Genetic aspects of pulmonary responses to inhaled pollutants. *Exp Toxicol Pathol* 2005; 57: 147–153.
- 27 Oudijk E-JD, Lammers J-WJ, Koenderman L. Systemic inflammation in chronic obstructive pulmonary disease. *Eur Respir J* 2003; 22: Suppl. 46, S5–S13.
- 28 Maritim AC, Sanders RA, Watkins JB 3rd. Diabetes, oxidative stress, and antioxidants: a review. *J Biochem Mol Toxicol* 2003; 17: 24–38.
- 29 Trasande L, Thurston GD. The role of air pollution in asthma and other pediatric morbidities. *J Allergy Clin Immunol* 2005; 115: 689–699.
- 30 Mayne ST. Antioxidant nutrients and chronic disease: use of biomarkers of exposure and oxidative stress status in epidemiologic research. *J Nutr* 2003; 133: Suppl. 3, 933S–940S.
- 31 Blanck HM, Bowman BA, Cooper GR, Myers GL, Miller DT. Laboratory issues: use of nutritional biomarkers. *J Nutr* 2003; 133: Suppl. 3, 888S–894S.
- 32 Handelman GJ. High-performance liquid chromatography analysis of cholesterol linoleate hydroperoxide in oxidized low density lipoproteins: calibration by conjugated diene internal standard. *Methods Enzymol* 1999; 300: 43–50.
- 33 Potischman N. Biologic and methodologic issues for nutritional biomarkers. *J Nutr* 2003; 133: Suppl. 3, 875S–880S.
- 34 Mickleborough TD, Rundell KW. Dietary polyunsaturated fatty acids in asthma- and exercise-induced bronchoconstriction. *Eur J Clin Nutr* 2005; 59: 1335–1346.
- 35 Costa DL, Dreher KL. Bioavailable transition metals in particulate matter mediate cardiopulmonary injury in healthy and compromised animal models. *Environ Health Perspect* 1997; 105: 1053–1060.
- 36 Soukup JM, Becker S. Human alveolar macrophage responses to air pollution particulates are associated with insoluble components of coarse material, including particulate endotoxin. *Toxicol Appl Pharmacol* 2001; 171: 20–26.
- 37 Gonzalez-Flecha B. Oxidant mechanisms in response to ambient air particles. *Mol Aspects Med* 2004; 25: 169–182.
- 38 Risom L, Moller P, Loft S. Oxidative stress-induced DNA damage by particulate air pollution. *Mutat Res* 2005; 592: 119–137.
- 39 Smith AD, Cowan JO, Brassett KP, Herbison GP, Taylor DR. Use of exhaled nitric oxide measurements to guide treatment in chronic asthma. *N Engl J Med* 2005; 35: 2163–2173.
- 40 Nel AE, Diaz-Sanchez D, Ng D, Hiura T, Saxon A. Enhancement of allergic inflammation by the interaction between diesel exhaust particles and the immune system. *J Allergy Clin Immunol* 1998; 102: 539–554.
- 41 Lim HB, Ichinose T, Miyabara Y, et al. Involvement of superoxide and nitric oxide on airway inflammation and hyperresponsiveness induced by diesel exhaust particles in mice. *Free Radic Biol Med* 1998; 25: 635–644.
- 42 Li N, Sioutas C, Cho A, et al. Ultrafine particulate pollutants induce oxidative stress and mitochondrial damage. *Environ Health Perspect* 2003; 111: 455–460.
- 43 Jimenez LA, Thompson J, Brown DA, et al. Activation of NF- κ B by PM(10) occurs via an iron-mediated mechanism in the absence of I κ B degradation. *Toxicol Appl Pharmacol* 2000; 166: 101–110.
- 44 Shukla A, Timblin C, Berube K, et al. Inhaled particulate matter causes expression of nuclear factor (NF)- κ B-related genes and oxidant-dependent NF- κ B activation *in vitro*. *Am J Respir Cell Mol Biol* 2000; 23: 182–187.
- 45 Aganasur K, Jeff Inmon P, Dailey LA, Madden MC, Ghio AJ, Gallagher JE. Air pollution particles mediated oxidative DNA base damage in a cell free system and in human airway epithelial cells in relation to particulate metal content and bioreactivity. *Chem Res Toxicol* 2001; 14: 879–887.
- 46 Squadrito GL, Cueto R, Dellinger B, Pryor WA. Quinoid redox cycling as a mechanism for sustained free radical generation by inhaled airborne particulate matter. *Free Radical Biol Med* 2001; 31: 1132–1138.
- 47 Hatzis C, Godleski JJ, Gonzalez-Flecha B, Wolfson JM, Koutrakis P. Ambient particulate matter exhibits direct inhibitory effects on oxidative stress enzymes. *Environ Sci Technol* 2006; 40: 2805–2811.
- 48 Takizawa H. Diesel exhaust particles and their effect on induced cytokine expression in human bronchial epithelial cells. *Curr Opin Allergy Clin Immunol* 2004; 4: 355–359.
- 49 Stenfors N, Nordenhall C, Salvi SS, et al. Different airway inflammatory responses in asthmatic and healthy humans exposed to diesel. *Eur Respir J* 2004; 23: 82–86.
- 50 Behndig AF, Mudway IS, Brown JL, et al. Airway antioxidant and inflammatory responses to diesel exhaust exposure in healthy humans. *Eur Respir J* 2006; 27: 359–365.
- 51 McDonald JD, Harrod KS, Seagrave J, Seilkop SK, Mauderly JL. Effects of low sulfur fuel and a catalyzed particle trap on the composition and toxicity of diesel emissions. *Environ Health Perspect* 2004; 112: 1307–1312.
- 52 Sun Q, Wang A, Jin X, et al. Long-term air pollution exposure and acceleration of atherosclerosis and vascular

- inflammation in an animal model. *JAMA* 2005; 294: 3003–3010.
- 53 Foucaud L, Bennisroune A, Klestadt D, Laval-Gilly P, Falla J. Oxidative stress induction by short time exposure to ozone on THP-1 cells. *Toxicol In Vitro* 2006; 20: 101–108.
 - 54 Janssen-Heininger YMW, Persinger RL, Korn SH, et al. Reactive nitrogen species and cell signaling. Implications for death or survival of lung epithelium. *Am J Respir Crit Care Med* 2002; 166: Suppl. 12, S9–S16.
 - 55 Kelly FJ, Tetley TD. Nitrogen dioxide depletes uric acid and ascorbic acid but not glutathione from lung lining fluid. *Biochem J* 1997; 325: 95–99.
 - 56 Persinger RL, Poynter ME, Ckless K, Janssen-Heininger YM. Molecular mechanisms of nitrogen dioxide induced epithelial injury in the lung. *Mol Cell Biochem* 2002; 234–235: 71–80.
 - 57 Xiao GG, Wang M, Li N, Loo JA, Nel AE. Use of proteomics to demonstrate a hierarchical oxidative stress response to diesel exhaust particle chemicals in a macrophage cell line. *J Biol Chem* 2003; 278: 50781–50790.
 - 58 Kelly FJ. Dietary antioxidants and environmental stress. *Procc Nutr Soc* 2004; 63: 579–585.
 - 59 Kelly FJ, Mudway IS. Protein oxidation at the air-lung interface. *Amino Acids* 2003; 25: 375–396.
 - 60 Freed AN, Cueto R, Pryor WA. Antioxidant transport modulates peripheral airway reactivity and inflammation during ozone exposure. *J Appl Physiol* 1999; 87: 7595–7603.
 - 61 Levine M, Katz A, Padayatt SJ. Vitamin C. In: Shils M, Shike M, Ross AC, et al., eds. *Modern Nutrition in Health and Disease*. Philadelphia, Lippincott Williams & Willkins, 2006; pp. 507–524.
 - 62 Mustacich D, Powis G. Thioredoxin reductase. *Biochem J* 2000; 346: 1–8.
 - 63 McCay PB. Vitamin E: interaction with free radicals and ascorbate. *Ann Rev Nutr* 1985; 5: 323–340.
 - 64 Burton GW, Ingold KU. Autooxidation of biological molecules. 1. The antioxidant activity of vitamin E and related chain-breaking phenolic antioxidants *in vitro*. *J Am Chem Soc* 1981; 103: 6472–6477.
 - 65 Traber MG. Vitamin E. In: Shils ME, Shike M, Ross AC, et al., eds. *Modern Nutrition in Health and Disease*. Philadelphia, Lippincott Williams & Willkins, 2006; pp. 396–311.
 - 66 Kolleck I, Sinha P, Rustow B. Vitamin E as an antioxidant of the lung: mechanisms of vitamin E delivery to alveolar type II cells. *Am J Respir Crit Care Med* 2002; 166: S62–S66.
 - 67 Linder MC. Nutrition and metabolism of trace elements. In: Linder MC, ed. *Nutrition Biochemistry and Metabolism with Clinical Application*. Norwalk, Appleton and Lange, 1991; pp. 213–276.
 - 68 Siems W, Wiswedel I, Salerno C, et al. β -Carotene breakdown products may impair mitochondrial functions—potential side effects of high-dose β -carotene supplementation. *J Nutr Biochem* 2005; 16: 385–397.
 - 69 Palozza P, Serini S, Di Nicuolo F, Piccioni E, Calviello G. Prooxidant effects of β -carotene in cultured cells. *Mol Aspects Med* 2003; 24: 353–362.
 - 70 Sies H, ed. *Oxidative Stress: Oxidants and Antioxidants*. San Diego, Academic Press, 1991.
 - 71 Burk RF, Levander OA. Selenium. In: Shils M, Olson JA, Shike M, Ross AC, eds. *Modern Nutrition in Health and Disease*. Baltimore, Williams, & Wilkins, 1999; pp. 265–276.
 - 72 Schwartz J, Weiss ST. Dietary factors and their relation to respiratory symptoms. *Am J Epidemiol* 1990; 132: 67–76.
 - 73 Rossi AG, Haslett C. Inflammation, cell injury, and apoptosis. In: Sais SE, ed. *Proinflammatory and Anti-inflammatory Peptides*. New York, Marcel Dekker, 1998.
 - 74 Holguin F, Tellez-Rojo MM, Hernandez M, et al. Air pollution and heart rate variability among the elderly in Mexico City. *Epidemiology* 2003; 14: 521–527.
 - 75 Gold DR, Litonjua A, Schwartz J, et al. Ambient pollution and heart rate variability. *Circulation* 2000; 101: 1267–1273.
 - 76 Zipes DP, Wellens HJ. Sudden cardiac death. *Circulation* 1998; 98: 2334–2351.
 - 77 Godleski JJ, Verrier RL, Koutrakis P, et al. Mechanisms of morbidity and mortality from exposure to ambient air particles. *Res Rep Health Eff Inst* 2000; 5–88.
 - 78 Katan MB, Deslypere JP, Van Birgelen AP, Pernders M, Zegwaard M. Kinetic of the incorporation of dietary fatty acids into serum cholesterly esters, erythrocyte membranes, and adipose tissue: an 18-month controlled study. *J Lipid Res* 1997; 38: 2012–2022.
 - 79 Kris-Etherton PM, Harris WS, Appel LJ, American Heart Association Nutrition Committee. Omega-3 fatty acids and cardiovascular disease: new recommendations from the American Heart Association. *Arterioscler Thromb Vasc Biol* 2003; 23: 151–152.
 - 80 Grievink L, Smit HA, Brunekreef B. Anti-oxidants and air pollution in relation to indicators of asthma and COPD: a review of the current evidence. *Clin Exp Allergy* 2000; 30: 1344–1354.
 - 81 Elsayed NM. Antioxidant mobilization in response to oxidative stress: a dynamic environmental-nutritional interaction. *Nutrition* 2001; 17: 828–834.
 - 82 Menzel DB. The toxicity of air pollution in experimental animals and humans: the role of oxidative stress. *Toxicol Lett* 1994; 72: 269–277.
 - 83 Menzel DB. Antioxidants in lung disease. *Toxicol Ind Health* 1993; 9: 323–336.
 - 84 Dunster C, Kelly FJ. Dietary modulation of lung epithelial lining fluid vitamin C concentration. *Respir Med* 1994; 88: 815.
 - 85 Pinho RA, Silveira PC, Silva LA, Luiz Streck E, Dal-Pizzol F, Moreira JC. N-acetylcysteine and deferoxamine reduce pulmonary oxidative stress and inflammation in rats after coal dust exposure. *Environ Res* 2005; 99: 355–360.
 - 86 Rhoden CR, Wellenius GA, Ghelfi E, Lawrence J, Gonzalez-Flecha B. PM-induced cardiac oxidative stress and dysfunction are mediated by autonomic stimulation. *Biochim Biophys Acta* 2005; 1725: 305–313.
 - 87 Mohsenin V. Effect of vitamin C on NO₂-induced airway hyperresponsiveness in normal subjects. A randomized double-blind experiment. *Am Rev Respir Dis* 1987; 136: 1408–1411.
 - 88 Chatham MD, Eppler Jr JH, Sauder LR, Green D, Kulle TJ. Evaluation of the effects of vitamin C on ozone-induced bronchoconstriction in normal subjects. *Ann N Y Acad Sci* 1987; 498: 269–279.
 - 89 Hackney JD, Linn WS, Buckley RD, et al. Vitamin E supplementation and respiratory effects of ozone in humans. *J Toxicol Environ Health* 1981; 7: 383–390.

- 90 Trenga CA, Williams PV, Koenig JQ. Dietary antioxidants attenuate ozone-induced bronchial hyperresponsiveness (BHR) in asthmatic adults. *Am J Resp Crit Care Med* 1997; 155: A732.
- 91 Samet JM, Hatch GE, Horstman D, et al. Effect of antioxidant supplementation on ozone-induced lung injury in human subjects. *Am J Respir Crit Care Med* 2001; 164: 819–825.
- 92 Mudway IS, Behndig AF, Helleday R, et al. Vitamin supplementation does not protect against symptoms in ozone-responsive subjects. *Free Radic Biol Med* 2006; 40: 1702–1712.
- 93 Grievink L, Zijlstra AG, Ke X, Brunekreef B. Acute effects of ozone on pulmonary function in antioxidant supplemented cyclists. *Eur Resp J* 1997; 10: Suppl. 25, 229S.
- 94 Grievink L, Zijlstra AG, Ke X, Brunekreef B. Double-blind intervention trial on modulation of ozone effects on pulmonary function by antioxidant supplements. *Am J Epidemiol* 1999; 149: 306–314.
- 95 Romieu I, Meneses F, Ramirez M, et al. Antioxidants supplementation and respiratory functions among workers exposed to high levels of ozone. *Am J Resp Crit Care Med* 1998; 158: 226–232.
- 96 Romieu I, Sierra-Monge JJ, Ramirez-Aguilar M, et al. Antioxidants supplementation and lung function among asthmatic children exposed to high levels of air pollutants. *Am J Respir Crit Care Med* 2002; 166: 703–709.
- 97 Romieu I, Sierra-Monge JJ, Ramirez-Aguilar M, et al. Genetic polymorphism of GSTM1 and antioxidant supplementation influence lung function in relation to ozone exposure in asthmatic children in Mexico City. *Thorax* 2004; 59: 8–10.
- 98 Sierra-Monge JJ, Ramirez-Aguilar M, Moreno-Macias H, et al. Antioxidant supplementation and nasal inflammatory responses among young asthmatics exposed to high levels of ozone. *Clin Exp Immunol* 2004; 138: 317–322.
- 99 Mickleborough TD. Dietary omega-3 polyunsaturated fatty acid supplementation and airway hyperresponsiveness in asthma. *J Asthma* 2005; 42: 305–314.
- 100 Dry J, Vincent D. Effect of a fish oil diet on asthma: results of a 1-year double-blind study. *Int Arch Allergy Appl Immunol* 1991; 95: 156–157.
- 101 Mickleborough TD, Murray RL, Ionescu AA, Lindley MR. Fish oil supplementation reduces severity of exercise-induced bronchoconstriction in elite athletes. *Am J Respir Crit Care Med* 2003; 168: 1181–1189.
- 102 Hodge L, Salome CM, Hughes JM, et al. Effect of dietary intake of omega-3 and omega-6 fatty acids on severity of asthma in children. *Eur Resp J* 1998; 11: 361–365.
- 103 Arm JP, Horton CE, Spur BW, Mencia-Huerta JM, Lee TH. The effects of dietary supplementation with fish oil lipids on the airways response to inhaled allergen in bronchial asthma. *Am Rev Respir Dis* 1989; 139: 1395–1300.
- 104 Thien FC, Mencia-Huerta JM, Lee TH. Dietary fish oil effects on seasonal hay fever and asthma in pollen-sensitive subjects. *Am Rev Respir Dis* 1993; 147: 1138–1143.
- 105 Marchioli R, Barzi F, Bomba E, et al. Early protection against sudden death by n-3 polyunsaturated fatty acids after myocardial infarction: time-course analysis of the results of the Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico (GISSI)-Prevenzione. *Circulation* 2002; 105: 1897–1903.
- 106 Romieu I, Tellez-Rojo MM, Lazo M, et al. Omega-3 fatty acid prevents heart rate variability reductions associated with particulate matter. *Am J Respir Crit Care Med* 2005; 172: 1534–1540.
- 107 Wayne EC. Cardiopulmonary health effects of air pollution: is a mechanism emerging? *Am J Respir Crit Care Med* 2005; 172: 1482–1484.
- 108 Schwartz J, Park SK, O'Neill MS, et al. Glutathione-S-transferase M1, obesity, statins, and autonomic effects of particles: gene-by-drug-by-environment interaction. *Am J Respir Crit Care Med* 2005; 172: 1529–1533.
- 109 Mudway I, Blomberg A, Helleday R, Frew A, Sandstrom T, Kelly FJ. Supplementation with vitamin C does not influence lung function in healthy subjects. *Eur Resp J* 2000; 16: Suppl. 31, 116S.
- 110 Paredi P, Kharitonov SA, Barnes PJ. Analysis of expired air for oxidation products. *Am J Respir Crit Care Med* 2002; 166: Suppl. 12, S31–S37.
- 111 Ratnawati R, Thomas PS. Exhaled nitric oxide in paediatric asthma. *Chron Respir Dis* 2005; 2: 163–174.
- 112 Uchida K. Role of reactive aldehyde in cardiovascular diseases. *Free Radic Biol Med* 2000; 28: 1685–1696.
- 113 Mannino DM. Epidemiology and global impact of chronic obstructive pulmonary disease. *Semin Respir Crit Care Med* 2005; 26: 204–210.
- 114 O'Neill MS, Veves A, Zanobetti A, et al. Diabetes enhances vulnerability to particulate air pollution-associated impairment in vascular reactivity and endothelial function. *Circulation* 2005; 111: 2913–2920.
- 115 Bateson TF, Schwartz J. Who is sensitive to the effects of particulate air pollution on mortality? A case-crossover analysis of effect modifiers. *Epidemiology* 2004; 15: 143–149.
- 116 Alberg A. The influence of cigarette smoking on circulating concentrations of antioxidant micronutrients. *Toxicology* 2002; 180: 121–137.
- 117 Dietrich M, Block G, Norkus EP, et al. Smoking and exposure to environmental tobacco smoke decrease some plasma antioxidants and increase γ -tocopherol *in vivo* after adjustment for dietary antioxidant intakes. *Am J Clin Nutr* 2003; 77: 160–166.
- 118 Hayes JD, Strange RC. Glutathione S-transferase polymorphism and their biological consequences. *Pharmacology* 2000; 61: 154–166.
- 119 Rushmore TH, Pickett CB. Glutathione S-transferase, structure, regulation, and therapeutic implications. *J Biol Chem* 1993; 268: 11475–11478.
- 120 Rebbeck TR. Molecular epidemiology of the human glutathione S-transferase genotype GSTM1 and GSTT1 in cancer susceptibility. *Cancer Epidemiol Biom Prev* 1997; 6: 733–743.
- 121 Gilliland FD, Lin Y, Dubeau L, et al. Effects of glutathione S-transferase M1, maternal smoking during pregnancy and environmental tobacco smoke on asthma and wheezing in children. *Am J Respir Crit Care Med* 2002; 166: 457–463.
- 122 Gilliland FD, Guaderman J, Vora H, Rappaport E, Dubeau L. Effects of glutathione S-transferase M1, T1, and P1 on childhood lung function growth. *Am J Respir Crit Care Med* 2002; 166: 710–716.

- 123** He JQ, Ruan J, Connett JE, Anthonisen NR, Paré PD, Sandford AJ. Antioxidant gene polymorphisms and susceptibility to a rapid decline in lung function in smokers. *Am J Respir Crit Care Med* 2002; 166: 323–328.
- 124** Kunzli N, Avol E, Gauderman J, et al. GSTM1 status modifies the effects of wildfire smoke on asthma symptoms. International Conference of the American Thoracic Society. *Am J Respir Crit Care Med* 2006: A502.
- 125** Corradi M, Alinovi R, Goldoni M, et al. Biomarkers of oxidative stress after controlled human exposure to ozone. *Toxicol Lett* 2002; 134: 219–225.
- 126** Gilliland FD, Li YF, Saxon A, Diaz-Sanchez D. Effect of glutathione-S-transferase M1 and P1 genotypes on xenobiotic enhancement of allergic responses: randomised, placebo-controlled crossover study. *Lancet* 2004; 363: 119–125.
- 127** Kopp EB, Medzhitov R. The Toll-receptor family and control of innate immunity. *Curr Opin Immunol* 1999; 11: 13–18.
- 128** Yang IA, Holz O, Jorres RA, et al. Association of tumor necrosis factor- α polymorphisms and ozone-induced change in lung function. *Am J Respir Crit Care Med* 2005; 171: 171–176.
- 129** Bayley JP, Ottenhoff TH, Verweij CL. Is there a future for TNF promoter polymorphisms? *Genes Immun* 2004; 5: 315–329.
- 130** Kleeberger SR, Levitt RC, Zhang LY, et al. Linkage analysis of susceptibility to ozone-induced lung inflammation in inbred mice. *Nat Genet* 1997; 17: 475–478.
- 131** Li H, Romieu I, Sienra-Monge JJ, et al. Genetic polymorphisms in arginase I and II and childhood asthma and atopy. *J Allergy Clin Immunol* 2006; 117: 119–126.
- 132** Bergamaschi E, De Palma G, Mozzoni P, et al. Polymorphism of quinone-metabolizing enzymes and susceptibility to ozone-induced acute effects. *Am J Respir Crit Care Med* 2001; 163: 1426–1431.
- 133** David GL, Romieu I, Sienra-Monge JJ, et al. Nicotinamide adenine dinucleotide (phosphate) reduced:quinone oxidoreductase and glutathione S-transferase M1 polymorphisms and childhood asthma. *Am J Respir Crit Care Med* 2003; 168: 1199–1104.
- 134** Tager IB, Weiss ST, Muñoz A, Rosner B, Speizer FE. Longitudinal study of the effects of maternal smoking on pulmonary function in children. *N Engl J Med* 1983; 309: 699–603.
- 135** Hanrahan JP, Tager IB, Segal MR, et al. The effect of maternal smoking during pregnancy on early infant lung function. *Am Rev Respir Dis* 1992; 145: 1129–1135.
- 136** Paige RC, Royce FH, Plopper CG, Buckpitt AR. Long-term exposure to ozone increases acute pulmonary centriacinar injury by 1-nitronaphthalene: I. Region-specific enzyme activity. *J Pharmacol Exp Ther* 2000; 295: 934–941.
- 137** Künzli N, Götschi T, Mudway IS, et al. Comparison of oxidative properties, light absorbance, total and elemental mass concentration of ambient PM_{2.5} collected at 20 European sites. *Environ Health Perspect* 2006; 114: 684–690.
- 138** Cadet J, Douki T, Ravanat JL. Artifacts associated with the measurement of oxidized DNA bases. *Environ Health Perspect* 1997; 105: 1034–1039.
- 139** Handelman GJ, Pryor WA. Evaluation of antioxidant status in humans. In: Papas AM, ed. Antioxidant status, diet, nutrition and health. Florida, CRC Press, Boca Raton, 1999; pp. 37–62.
- 140** Ghiselli A, Serafini M, Natella F, Scaccini C. Total antioxidant capacity as a tool to assess redox status: critical view and experimental data. *Free Radic Biol Med* 2000; 30: 1036–1044.
- 141** Huang D, Ou B, Prior RL. The chemistry behind antioxidant capacity assays. *J Agric Food Chem* 2005; 53: 1841–1856.
- 142** Wayner DD, Burton GW, Ingold KU, Locke S. Quantitative measurement of the total, peroxy radical-trapping antioxidant capability of human blood plasma by controlled peroxidation. The important contribution made by plasma proteins. *FEBS Lett* 1985; 187: 33–37.
- 143** Mayne ST. Antioxidant nutrients and chronic disease: use of biomarkers of exposure and oxidative stress status in epidemiologic research. *J Nutr* 2003; 133: Suppl. 3, 933S–940S.
- 144** Cherubini A, Ruggiero C, Polidori MC, Mecocci P. Potential markers of oxidative stress in stroke. *Free Radic Biol Med* 2005; 39: 841–852.
- 145** Del Rio D, Stewart AJ, Pellegrini N. A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutr Metab Cardiovasc Dis* 2005; 15: 316–328.
- 146** Karatas F, Karatepe M, Baysar A. Determination of free malondialdehyde in human serum by high-performance liquid chromatography. *Anal Biochem* 2002; 311: 76–79.
- 147** Zarkovic N. 4-hydroxynonenal as a bioactive marker of pathophysiological processes. *Mol Aspects Med* 2003; 24: 281–291.
- 148** Gueraud F, Peiro G, Bernard H, et al. Enzyme immunoassay for a urinary metabolite of 4-hydroxynonenal as a marker of lipid peroxidation. *Free Radic Biol Med* 2006; 40: 54–62.
- 149** Paredi P, Kharitonov SA, Barnes PJ. Elevation of exhaled ethane concentration in asthma. *Am J Respir Crit Care Med* 2000; 162: 1450–1454.
- 150** Meagher EA, FitzGerald GA. Indices of lipid peroxidation *in vivo*: strengths and limitations. *Free Radic Biol Med* 2000; 28: 1745–1750.
- 151** Kanoh S, Kobayashi H, Motoyoshi K. Exhaled ethane: an *in vivo* biomarker of lipid peroxidation in interstitial lung diseases. *Chest* 2005; 128: 2387–2392.
- 152** Wood LG, Gibson PG, Garg ML. Biomarkers of lipid peroxidation, airway inflammation and asthma. *Eur Respir J* 2003; 21: 177–186.
- 153** Griffiths HR, Moller L, Bartosz G, et al. Biomarkers. *Mol Aspects Med* 2002; 23: 101–208.
- 154** Puhl H, Waeg G, Esterbauer H. Methods to determine oxidation of low-density lipoproteins. *Methods Enzymol* 1994; 233: 425–441.
- 155** Cole TG, Parikh N. High-throughput measurement of oxidizability of low-density lipoproteins suitable for use in clinical trials. *Clin Chem* 1999; 45: 696–699.
- 156** Ahotupa M, Asankari TJ. Baseline diene conjugation in LDL lipids: an indicator of circulating oxidized LDL. *Free Radic Biol Med* 1999; 27: 1141–1150.

- 157** Uno M, Kitazato KT, Nishi K, Itabe H, Nagahiro S. Raised plasma oxidised LDL in acute cerebral infarction. *J Neurol Neurosurg Psychiatry* 2003; 74: 312–316.
- 158** Polidori MC, Frei B, Cherubini A, et al. Increased plasma levels of lipid hydroperoxides in patients with ischemic stroke. *Free Radic Biol Med* 1998; 25: 561–567.
- 159** Rosias P, Robroeks C, Hendriks J, Dompeling E, Jobsis Q. Exhaled breath condensate: a space odyssey, where no one has gone before. *Eur Respir J* 2004; 24: 189–190.
- 160** Montuschi P, Barnes PJ. Exhaled leukotrienes and prostaglandins in asthma. *J Allergy Clin Immunol* 2002; 109: 615–620.
- 161** Montuschi P, Ragazzoni E, Valente S, et al. Validation of eicosanoid measurements in exhaled breath condensate. *Eur Respir J* 2002; 20: Suppl. 38, 422s–423s.
- 162** Montuschi P, Ragazzoni E, Valente S, et al. Validation of 8-isoprostane and prostaglandin E(2) measurements in exhaled breath condensate. *Inflamm Res* 2003; 52: 502–507.
- 163** Latzin P, Griese M. Exhaled hydrogen peroxide, nitrite and nitric oxide in healthy children: decrease of hydrogen peroxide by atmospheric nitric oxide. *Eur J Med Res* 2002; 7: 353–358.
- 164** Tsikas D. Simultaneous derivatization and quantification of the nitric oxide metabolites nitrite and nitrate in biological fluids by gas chromatography/mass spectrometry. *Anal Chem* 2000; 72: 4046–4072.
- 165** Corradi M, Folesani G, Andreoli R, et al. Aldehydes and glutathione in exhaled breath condensate of children with asthma exacerbation. *Am J Respir Crit Care Med* 2003; 167: 395–399.
- 166** Dziedzic B, Mazanowska-Gajdowicz J, Walczewska A, Nowak D. Application of Griess method for NO₂⁻/NO₃⁻ measurement in expired breath condensate. *Eur Respir J* 2002; 20: Suppl. 38, 92s.
- 167** Hensley K, Williamson KS, Floyd RA. Measurement of 3-nitrotyrosine and 5-nitro- γ -tocopherol by high-performance liquid chromatography with electrochemical detection. *Free Radic Biol Med* 2000; 28: 520–528.
- 168** Wink DA, Kim S, Coffin D, et al. Detection of S-nitrosothiols by fluorometric and colorimetric methods. *Methods Enzymol* 1999; 301: 201–211.
- 169** Larstad M, Soderling AS, Olin AC, Caidahl K, Toren K. Mass-selective determination of free 3-nitrotyrosine in breath condensate. *Eur Respir J* 2002; 20: Suppl. 31, 484s.
- 170** Halliwell B. Why and how should we measure oxidative DNA damage in nutritional studies? How far have we come? *Am J Clin Nutr* 2000 Nov, 72: 1082–1087.
- 171** Santella RM. Immunological methods for detection of carcinogen-DNA damage in humans. *Cancer Epidemiol Biomarkers Prev* 1999; 8: 733–739.
- 172** Long L, McCabe DR, Dolan ME. Determination of 8-oxoguanine in human plasma and urine by high-performance liquid chromatography with electrochemical detection. *J Chromatogr B Biomed Sci Appl* 1999; 731: 241–249.
- 173** Sauvaigo S, Petec-Calin C, Caillat S, Odin F, Cadet J. Comet assay coupled to repair enzymes for the detection of oxidative damage to DNA induced by low doses of γ -radiation: use of YOYO-1, low-background slides, and optimized electrophoresis conditions. *Anal Biochem* 2002; 303: 107–109.
- 174** Collins AR. Assays for oxidative stress and antioxidant status: applications to research into the biological effectiveness of polyphenols. *Am J Clin Nutr* 2005; 81: Suppl. 1, 261S–267S.
- 175** Gedik CM, Boyle SP, Wood SG, Vaughan NJ, Collins AR. Oxidative stress in humans: validation of biomarkers of DNA damage. *Carcinogenesis* 2002; 23: 1441–1446.
- 176** Collins AR, Duthie SJ, Dobson VL. Direct enzymic detection of endogenous oxidative base damage in human lymphocyte DNA. *Carcinogenesis* 1993; 14: 1733–1735.
- 177** Hu JJ, Chi CX, Frenkel K, et al. α -Tocopherol dietary supplement decreases titers of antibody against 5-hydroxymethyl-2'-deoxyuridine (HMdU). *Cancer Epidemiol Biomarkers Prev* 1999; 8: 693–698.
- 178** Frenkel K, Karkoszka J, Glassman T, et al. Serum autoantibodies recognizing 5-hydroxymethyl-2'-deoxyuridine, an oxidized DNA base, as biomarkers of cancer risk in women. *Cancer Epidemiol Biomarkers Prev* 1998; 7: 49–57.
- 179** Buss H, Chan TP, Sluis KB, Domigan NM, Winterbourn CC. Protein carbonyl measurement by a sensitive ELISA method. *Free Radic Biol Med* 1997; 23: 361–366.
- 180** Corradi M, Saglia S, Majori M, Zanini A, Pesci A, Cuomo A. A new technique for measurement of nitrate in breath condensate. *Am J Respir Crit Care Med* 2000; 161: A395.
- 181** Beeh KM, Beier J, Haas IC, Kornmann O, Micke P, Buhl R. Glutathione deficiency of the lower respiratory tract in patients with idiopathic pulmonary fibrosis. *Eur Respir J* 2002; 19: 1119–1123.
- 182** Griese M, Ramakers J, Krasselt A, et al. Improvement of alveolar glutathione and lung function but not oxidative state in cystic fibrosis. *Am J Respir Crit Care Med* 2004; 169: 822–828.
- 183** Cereser C, Guichard J, Drai J, et al. Quantification of reduced and total glutathione at the femtomole level by high-performance liquid chromatography with fluorescence detection: application to red blood cells and cultured fibroblast. *J Chromatogr B Biomed Sci Appl* 2001; 752: 123–132.
- 184** Kuhn KS, Krasselt AI, Furst P. Glutathione and glutathione metabolites in small tissue samples and mucosal biopsies. *Clin Chem* 2000; 46: 1003–1005.
- 185** James SJ, Cutler P, Melnyk S, et al. Metabolic biomarkers of increased oxidative stress and impaired methylation capacity in children with autism. *Am J Clin Nutr* 2004; 80: 1611–1617.
- 186** Schleiss MB, Holz O, Behnke M, Richter K, Magnussen H, Jorres RA. The concentration of hydrogen peroxide in exhaled air depends on expiratory flow rate. *Eur Respir J* 2000; 16: 1115–1118.
- 187** Horvath I, MacNee W, Kelly FJ, et al. Haemoxygenase-1 induction and exhaled markers of oxidative stress in lung diseases. *Eur Respir J* 2001; 18: 420–430.
- 188** van Beurden WJ, Dekhuijzen PN, Smeenk FW. Exhaled biomarkers in COPD: their potential role in diagnosis, treatment and prognosis. *Monaldi Arch Chest Dis* 2002; 57: 258–267.
- 189** Broeckaert F, Arsalane K, Hermans C, et al. Serum clara cell protein: a sensitive biomarker of increased lung

- epithelium permeability caused by ambient ozone. *Environ Health Perspect* 200; 108: 533–537.
- 190** Bernard A, Crbonnelle S, Nickmilder M, de Burbure C. Non-invasive biomarkers of pulmonary damage and inflammation: application to children exposed to ozone and trichlormine. *Toxicol Appl Pharmacol* 2005; 206: 185–190.
- 191** Hermans C, Aly O, Nyberg BI, Peterson C, Bernard A. Determinants of clara cell protein (CC16) concentration in serum: a reassessment with two different immunoassays. *Clin Chim Acta* 1998; 272: 101–110.
- 192** Blomberg A, Mudway I, Svensson M, *et al.* Clara cell protein as a biomarker for ozone-induced lung injury in humans. *Eur Respir J* 2003; 22: 883–888.
- 193** Miwa K, Kishimoto C, Nakamura H, *et al.* Serum thioredoxin and α -tocopherol concentrations in patients with major risk factors. *Circ J* 2005; 69: 291–294.
- 194** Nakamura H, Vaage J, Valen G, Padilla CA, Björnstedt M, Holmgren A. Measurements of plasma glutaredoxin and thioredoxin in healthy volunteers and during open-heart surgery. *Free Radic Biol Med* 1998; 24: 1176–1186.
- 195** Hokamaki J, Kawano H, Soejima H, *et al.* Plasma thioredoxin levels in patients with unstable angina. *Int J Cardiol* 2005; 99: 225–231.
- 196** Khachik F, Spangler CJ, Smith JC Jr, Canfield LM, Steck A, Pfander H. Identification, quantification, and relative concentrations of carotenoids and their metabolites in human milk and serum. *Anal Chem* 1997; 69: 1873–1881.
- 197** Stahl W, Sies H. Uptake of lycopene and its geometrical isomers is greater from heat-processed than from unprocessed tomato juice in humans. *J Nutr* 1992; 122: 2161–2166.
- 198** Wood LG, Garg ML, Blake RJ, Garcia-Caraballo S, Gibson PG. Airway and circulating levels of carotenoids in asthma and healthy controls. *J Am Coll Nutr* 2005; 24: 448–455.
- 199** Nierenberg DW, Peng YM, Alberts DS. Methods for determination of retinoids, α -tocopherols, and carotenoids in human serum, plasma and other tissues. In: Moon TE, Micozzi MS, eds. *Nutrition and Cancer Prevention*. Marcel Dekker, New York, 1989; pp. 181–209.
- 200** Margolis SA, Duester DL. Measurement of ascorbic acid in human plasma and serum: stability, intralaboratory repeatability, and interlaboratory reproducibility. *Clin Chem*. 1996; 42: 1257–1262.
- 201** Alegria A, Barbera R, Clemente G, Farre R, Garcia MJ, Lagarda MJ. Selenium and glutathione peroxidase reference values in whole blood and plasma of a reference population living in Valencia, Spain. *J Trace Elem Med Biol* 1996; 10: 223–228.
- 202** Hambidge M. Biomarkers of trace mineral intake and status. *J Nutr* 2003; 133: Suppl. 3, 948S–955S.
- 203** Muñoz-Olivas R, Donard OFX, Gilon N, Potin-Gautier M. Speciation of organic selenium compounds by high-performance liquid chromatography-inductively coupled plasma mass spectrometry in natural samples. *J Anal Atomic Spectr* 1996; 11: 1171–1176.
- 204** Koyama H, Omura K, Ejima A, Kasanuma Y, Watanabe C, Satoh H. Separation of selenium-containing proteins in human and mouse plasma using tandem high-performance liquid chromatography columns coupled with inductively coupled plasma-mass spectrometry. *Anal Biochem* 1999; 267: 84–91.
- 205** Dwyer JT, Peterson JJ. Measuring flavonoid intake: need for advanced tools. *Public Health Nutr* 2002; 5: 925–930.
- 206** Radtke J, Linseisen J, Wolfram G. Fasting plasma concentrations of selected flavonoids as markers of their ordinary dietary intake. *Eur J Nutr* 2002; 41: 203–209.
- 207** Lampe JW. Isoflavonoid and lignan phytoestrogens as dietary biomarkers. *J Nutr* 2003; 133: Suppl. 3, 956S–964S.
- 208** Adlercreutz H, Fotsis T, Bannwart C, Wahala K, Brunow G, Hase T. Isotope dilution gas chromatographic-mass spectrometric method for the determination of lignans and isoflavonoids in human urine, including identification of genistein. *Clin Chim Acta*. 1991; 199: 263–278.
- 209** Franke AA, Custer LJ, Cerna CM, Narala K. Rapid HPLC analysis of dietary phytoestrogens from legumes and from human urine. *Proc Soc Exp Biol Med* 1995; 208: 18–26.
- 210** Arab L. Biomarkers of fat and fatty acid intake. *J Nutr* 2003; 133: Suppl. 3, 925S–932S.
- 211** Kohlmeier L. Future of dietary exposure assessment. *Am J Clin Nutr* 1995; 61: Suppl. 3, 702S–709S.
- 212** Bates CJ, Thurnham DI, Bringham SA, Margatts BM, Nelson M. Biochemical markers of nutrient intake. In: Margetts BM, Nelson M, eds. *Design Concepts in Nutritional Epidemiology*. 2nd Edn. Oxford University Press, Oxford, 1997; pp. 170–240.
- 213** Mason JB. Biomarkers of nutrient exposure and status in one-carbon (methyl) metabolism. *J Nutr* 2003; 133: Suppl. 3, 941S–947S.
- 214** Bagley PJ, Selhub J. Analysis of folates using combined affinity and ion-pair chromatography. *Methods Enzymol* 1997; 281: 16–25.
- 215** Duggan C, MacLeod WB, Krebs NF, *et al.* Plasma zinc concentrations are depressed during the acute phase response in children with falciparum malaria. *J Nutr* 2005; 135: 802–807.
- 216** Batchet L, Vaja S, Treacher D, Kinerons M, Swaminathan R. Erythrocyte zinc in hospital patients. *Ann Clin Biochem* 2005; 42: 448–452.
- 217** Torrejon CS, Castillo-Duran C, Hertrampf ED, Ruz M. Zinc and iron nutrition in Chilean children fed fortified milk provided by the Complementary National Food Program. *Nutrition* 2004; 20: 177–180.
- 218** Smith JC Jr, Butrimovitz GP, Purdy WC. Direct measurement of zinc in plasma by atomic absorption spectroscopy. *Clin Chem*. 1979; 25: 1487–1491.
- 219** Pekkanen J, Sunyer J, Chinn S. Nondifferential disease misclassification may bias incidence risk ratios away from the null. *J Clin Epidemiol* 2006; 59: 281–289.
- 220** Pekkanen J, Sunyer J, Antó JM. Operational definitions of asthma in studies on its aetiology. *Eur Respir J* 2005; 26: 28–35.